

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, 15/60, A01H 5/00	A1	(11) International Publication Number: WO 96/30530 (43) International Publication Date: 3 October 1996 (03.10.96)
(21) International Application Number: PCT/US96/03635 (22) International Filing Date: 22 March 1996 (22.03.96) (30) Priority Data: 08/409,297 24 March 1995 (24.03.95) US (71) Applicant: PIONEER HI-BRED INTERNATIONAL, INC. [CA/US]; 700 Capital Square, 400 Locust Street, Des Moines, IO 50390 (US). (72) Inventors: BASZCZYNSKI, Chris, L.; 7305 Benton Drive, Urbandale, IO 50322 (US). BARBOUR, Eric; #28-5513 Aurora Avenue, Des Moines, IO 50310 (US). MIKI, Brian; 1876 Dorset Drive, Ottawa, Ontario K1H 5V1 (CA). (74) Agents: BENT, Stephen, A. et al.; Foley & Lardner, Suite 500, 3000 K Street, N.W., Washington, DC 20007-5109 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: ALS3 PROMOTER (57) Abstract The <i>Brassica</i> ALS3 promoter is operably linked to a foreign structural gene to provide high level, and generally constitutive or tissue general expression of the structural gene in transformed plants. The <i>Brassica</i> ALS3 promoter exhibits a non-tissue-preferred mode of expression at a level comparable to, and in some cases higher than, the widely used CaMV 35S promoter. Accordingly, DNA constructs comprising the <i>Brassica</i> ALS3 promoter operably linked to any number of different gene coding regions can be used for constitutive and tissue-general expression of the gene in transformed plants. The <i>Brassica</i> ALS3 promoter is used to direct expression of agronomically important genes and selectable marker genes.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

ALS3 PROMOTER

BACKGROUND OF THE INVENTION

I. Field of the Invention

5 The present invention relates to a novel ALS3 promoter which generally drives constitutive and generally non-tissue-preferred expression of operably linked foreign genes in transformed plants. In particular, this invention is directed to DNA constructs in which a *Brassica* ALS3 promoter is operably linked to
10 a foreign structural gene, and to using the DNA construct to produce, in a transformed plant, a protein which is encoded by the structural gene. The *Brassica* ALS3 promoter is used to direct expression of agronomically important genes and selectable marker genes.

15 II. Background

Acetolactate synthase (ALS), which is also known as acetohydroxy acid synthase (AHAS), catalyses the first step in the biosynthesis of the branched chain amino acids leucine, isoleucine and valine. It has also been
20 shown to be the site of action of sulfonylurea and imidazolinone based herbicides. See, for example, Chaleff, R.S. and C.J. Mauvais, *Science* 224:1443 (1984) and Shaner et al., *Plant Physiol.* 76:545 (1984). A number of different ALS genes from *Brassica napus* have
25 been cloned and characterized. See, for example Wiersma et al., *Mol. Gen. Genetics* 219:413 (1989) and Rutledge et al., loc. cit. 229:31 (1991).

Rutledge et al. (1991) reported that the *B. napus* rapeseed cultivar Topas contains an ALS multigene family
30 comprised of five genes. DNA sequence analysis of the structural genes revealed that the ALS1 and ALS3 genes shared extensive sequence homology. In contrast, the ALS2 gene has diverged significantly from the ALS1 and ALS3 genes and has unique features in the coding region
35 of the mature polypeptide, transit peptide and upstream non-coding region. The ALS2 gene therefore may encode a

polypeptide with a distinct function from that of ALS1 and ALS3. The ALS4 and ALS5 genes have interrupted coding regions and therefore may be defective.

Experiments conducted with the promoter—of the
5 *Arabidopsis thaliana* ALS gene revealed that the *A. thaliana* ALS promoter is significantly less effective in driving gene expression than the CaMV 35S promoter. Odell et al., *Plant Physiol.* 94(4):1647-1654 (1990) replaced the *A. thaliana* ALS promoter with the CaMV 35S
10 promoter and observed a 25-fold increase in the level of ALS mRNA accompanied by a 2-fold increase in ALS enzyme level and a 3-fold increase in sulfonylurea tolerance. These observations indicate that the ALS gene is regulated post-transcriptionally and that the *A. thaliana*
15 ALS promoter is significantly less effective in driving gene expression than the CaMV 35S promoter.

The number of isolated and characterized constitutive generally non-tissue-preferred plant promoters available for expression of foreign proteins in transgenic plants
20 is very limited. Well known examples of promoters with constitutive and tissue generated expression patterns include those associated with the CaMV 35S, *Agrobacterium* nopaline synthase, and maize ubiquitin genes. See Odell et al., *Plant Mol. Biol.* 10(3):263-272 (1988), Herrera-
25 Estrella et al., *Nature* 303:209-213 (1983) and Fox et al., *Va. J. Sci.* 43(2):287 (1992).

There is a critical need for a broader repertoire of strong constitutive and generally non-tissue-preferred plant promoters. A broader array of constitutive and
30 generally non-tissue-preferred plant promoters that are expressed at high levels, that is, that drive expression of operably linked genes at a level comparable to the CaMV 35S promoter, would allow the genetic engineer to analyze the relative strengths of the available promoters
35 and select promoters that provide the required level of expression of foreign genes in transformed plants. A

- 3 -

selected promoter might provide optimum levels of expression for the first gene but may be either too strong or too weak for use in driving the expression of a second gene. Consequently, additional constitutive and tissue general promoters are needed to optimize foreign gene expression in plants.

There is also a need for additional strong constitutive and generally non-tissue preferred promoters for construction of plants transformed with multiple foreign genes. Numerous difficulties have arisen when two or more different genes are introduced into a plant wherein each of the genes are operably linked to the same or similar promoters. Some of these difficulties include (1) gene inactivation; (2) recombination as a result of pairing along homologous regions within the nucleotide sequence of the promoter leading to cross-over events and loss of the intervening region prior, or subsequent to, integration; and (3) competition among different copies of the same promoter region for binding of promoter-specific transcription factors or other regulatory DNA-binding proteins. A need therefore exists for a broader repertoire of strongly constitutive and tissue general promoters to be used for expression of foreign genes in transformed plants.

SUMMARY OF THE INVENTION

Accordingly, it is an object of the present invention to provide a strong, constitutive promoter which can effect high level, generally non-tissue-preferred expression of an operably linked, foreign gene in transformed plants.

It is another object of the present invention to add to the limited repertoire of generally non-tissue-preferred promoters available for the transformation of plants with multiple genes.

In achieving these and other objects, there has been provided, in accordance with one aspect of the present

invention, an isolated DNA molecule comprising a promoter operably linked to a foreign structural gene, wherein said promoter comprises a nucleotide sequence corresponding to the sequence of a polynucleotide from the group consisting of (1) a *XbaI/NcoI* fragment 5-prime to the *Brassica napus* ALS3 structural gene or (2) a nucleotide sequence that has substantial sequence similarity with said *XbaI/NcoI* fragment. Other objects of the present invention include providing an isolated DNA molecule wherein the promoter is operably linked to an agronomically important gene or a selectable marker gene, an isolated DNA that is part of an expression vector and an expression vector carrying the isolated DNA molecule that is present in a transformed host.

It is another object of the present invention to provide an isolated DNA molecule comprising a promoter operably linked to a foreign structural gene, wherein said promoter comprises a nucleotide sequence corresponding to the sequence of a polynucleotide from the group consisting of (1) SEQ ID NO: 1 or (2) a nucleotide sequence that has substantial sequence similarity with SEQ ID NO: 1. Other objects of the present invention include providing an isolated DNA molecule wherein the promoter is operably linked to an agronomically important gene or a selectable marker gene, an isolated DNA that is part of an expression vector and an expression vector carrying the isolated DNA molecule that is present in a transformed host.

Another object of the present invention is to provide a method of using an ALS3 promoter to produce a foreign protein in a transformed host plant, comprising the steps of (1) constructing an expression vector comprising a promoter operably linked to a foreign structural gene, wherein the promoter comprises a nucleotide sequence corresponding to the sequence of a polynucleotide from the group consisting of (a) a *XbaI/NcoI* fragment 5-prime

- 5 -

to the *Brassica napus* ALS3 structural gene or (b) a nucleotide sequence that has substantial sequence similarity with said *XbaI/NcoI* fragment; and (2) transforming a host.

5 Yet another object of the present invention is to provide a method of using an ALS3 promoter to produce a foreign protein in a transformed host plant, comprising the steps of (1) constructing an expression vector comprising a promoter operably linked to a foreign
10 structural gene, wherein said promoter comprises a nucleotide sequence corresponding to the sequence of a polynucleotide from the group consisting of (a) SEQ ID NO: 1 or (b) a nucleotide sequence that has substantial sequence similarity with SEQ ID NO: 1; and (2)
15 transforming a host.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples,
20 while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

25 **BRIEF DESCRIPTION OF THE DRAWINGS**

Figure 1 presents the nucleotide sequence [SEQ ID NO: 1] of a 1063 bp *XbaI/NcoI* fragment that comprises the promoter 5-prime to the *B. napus* ALS3 wild type structural gene. The *XbaI* and *NcoI* restriction sites are
30 underlined. The ATG start codon is found within the *NcoI* restriction site.

Figure 2 presents a restriction map of a 1063 bp *XbaI/NcoI* fragment which comprises the *B. napus* ALS3 wild type structural gene.

35 Figure 3 presents a map of pPHI4960, a binary vector containing the *B. napus* ALS3 promoter (ALS3-Pro) driving

- 6 -

uidA (GUS) gene expression. Plasmid pPHI4960 was used to transform canola and sunflower. LB and RB represent the left and right borders, respectively, of the *Agrobacterium* Ti plasmid T-DNA region. The region between the left and right borders also includes a
5 CaMV 35S promoter (35S-Pro) driving expression of the NPTII gene for kanamycin selection of transformed plants.

Figure 4 presents a map of pPHI6333 which is a binary vector containing a cassette comprising the *B. napus* ALS3 promoter (BNALS3 PROM), NPTII gene and the 3-prime terminator sequence from the potato proteinase inhibitor gene (PINII). Plasmid pPHI6333 was used to transform sunflower to ascertain the suitability of this promoter for driving selectable marker expression. LB and RB
10 represent the left and right borders, respectively, of the *Agrobacterium* Ti plasmid T-DNA region.
15

Figure 5 presents a map of plasmid pPHI3042 which contains the NPTII and GUS structural genes operably linked to the CaMV 35S promoter.

Figure 6 presents a map of plasmid pPHI5765 which contains the NPTII structural gene operably linked to the CaMV 35S promoter (355PRO) and the potato proteinase inhibitor gene terminator sequence (PINII).
20

Figure 7 is a histogram showing GUS activity in plant seedlings transformed with expression cassettes in which the GUS gene is expressed from either the CaMV 35S or ALS3 promoters. GUS activity was measured in cotyledon, hypocotyl, meristem and root tissues.
25

Figure 8 is a histogram showing GUS activity in vegetative stage plants transformed with expression cassettes in which the GUS gene is expressed from either the CaMV 35S or ALS3 promoters. GUS activity was measured in leaf, petiole, meristem, stem and root tissues.
30

Figure 9 is a histogram showing GUS activity in flowering stage plants transformed with expression
35

- 7 -

cassettes in which the GUS gene is expressed from either the CaMV 35S or ALS3 promoters. GUS activity was measured in flower, leaf stem and root tissues.

5 Figure 10 is a histogram showing GUS activity in mature plants transformed with expression cassettes in which the GUS gene is expressed from either the CaMV 35S or ALS3 promoters. GUS activity was measured in leaf, stem, pod and seed tissues.

DETAILED DESCRIPTION

10 I. Definitions

In the description that follows, a number of terms are used extensively. The following definitions are provided to facilitate understanding of the invention.

15 A structural gene is a DNA sequence that is transcribed into messenger RNA (mRNA) which is then translated into a sequence of amino acids characteristic of a specific polypeptide.

A promoter is a DNA sequence that directs the transcription of a structural gene. Typically, a
20 promoter is located in the 5' region of a gene, proximal to the transcriptional start site of a structural gene. If a promoter is an inducible promoter, then the rate of transcription increases in response to an inducing agent. For example, a promoter may be regulated in a tissue-specific
25 manner such that it is only active in transcribing the associated coding region in a specific tissue type(s) such as leaves, roots or meristem.

In contrast, the rate of transcription is not generally regulated by an inducing agent if the promoter
30 is a constitutive promoter. The promoter may be tissue-general, also known as non-tissue-preferred, such that it is active in transcribing the associated coding region in a variety of different tissue types.

A core promoter contains essential nucleotide
35 sequences for promoter function, including the TATA box

and start of transcription. By this definition, a core promoter may or may not have detectable activity in the absence of specific sequences that may enhance the activity.

5 An isolated DNA molecule is a fragment of DNA that is not integrated in the genomic DNA of an organism. For example, the promoter of the ALS3 gene is a DNA fragment that has been separated from the genomic DNA of *Brassica napus*.

10 Complementary DNA (cDNA) is a single-stranded DNA molecule that is formed from an mRNA template by the enzyme reverse transcriptase. Typically, a primer complementary to portions of mRNA is employed for the initiation of reverse transcription. Those skilled in
15 the art also use the term "cDNA" to refer to a double-stranded DNA molecule consisting of such a single-stranded DNA molecule and its complementary DNA strand.

To operably link one nucleotide sequence to another refers to joining two heterologous DNA fragments to
20 produce a chimeric DNA construct that has biological activity. For example, an isolated DNA fragment comprising a promoter from a first gene, such as the ALS3 gene, is operably linked to an isolated DNA fragment comprising the structural gene from a second heterologous
25 gene. The resulting chimeric DNA construct is functional when the ALS3 promoter is shown to initiate transcription of the heterologous structural gene.

The term expression refers to the biosynthesis of a gene product. For example, in the case of a structural
30 gene, expression involves transcription of the structural gene into mRNA and the translation of mRNA into one or more polypeptides.

A cloning vector is a DNA molecule, such as a plasmid, cosmid, or bacteriophage, that has the
35 capability of replicating autonomously in a host cell. Cloning vectors typically contain one or a small number

of restriction endonuclease recognition sites at which foreign DNA sequences can be inserted in a determinable fashion without loss of an essential biological function of the vector, as well as a marker gene that is suitable
5 for use in the identification and selection of cells transformed with the cloning vector. Marker genes typically include genes that provide tetracycline resistance or ampicillin resistance.

An expression vector is a DNA molecule comprising a
10 gene that is expressed in a host cell. Typically, gene expression is placed under the control of certain regulatory elements, including constitutive or inducible promoters, tissue-specific regulatory elements, and enhancers. Such a gene is said to be "operably linked
15 to" the regulatory elements.

A foreign gene refers in the present description to a DNA sequence that is operably linked to at least one heterologous regulatory element. For example, any gene
20 other than the ALS3 structural gene is considered to be a foreign gene if the expression of that gene is controlled by the ALS3 promoter.

A recombinant host may be any prokaryotic or eukaryotic cell that contains either a cloning vector or expression vector. This term also includes those
25 prokaryotic or eukaryotic cells that have been genetically engineered to contain the cloned gene(s) in the chromosome or genome of the host cell.

A transgenic plant is a plant having one or more plant cells that contain an expression vector.

30 In eukaryotes, RNA polymerase II catalyzes the transcription of a structural gene to produce mRNA. A DNA molecule can be designed to contain an RNA polymerase II template in which the RNA transcript has a sequence that is complementary to that of a specific mRNA. The
35 RNA transcript is termed an antisense RNA and a DNA sequence that encodes the antisense RNA is termed an

- 10 -

antisense gene. Antisense RNA molecules are capable of binding to mRNA molecules, resulting in an inhibition of mRNA translation.

A first nucleotide sequence has substantial-sequence
5 similarity to the nucleotide sequence of Figure 1 [SEQ. ID NO: 1] if the former sequence share a similarity of at least 65% with the Figure 1 sequence and is a constitutive promoter active in directing the transcription of an operably linked foreign structural
10 gene in plants. Sequence similarity determinations can be performed, for example, using the FASTA program (Genetics Computer Group; Madison, WI). Alternatively, sequence similarity determinations can be performed using
15 BLASTP (Basic Local Alignment Search Tool) of the Experimental GENIFO® BLAST Network Service. See Altschul et al., *J. Mol. Biol.* 215:403 (1990). Also, see Pasternak et al., "Sequence Similarity Searches, Multiple Sequence Alignments, and Molecular Tree Building," in METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY,
20 Glick et al. (eds.), pages 251-267 (CRC Press 1993). Promoter activity of the isolated nucleotide sequence can be assayed by means of fusing the nucleotide sequence to a coding region of a foreign reporter gene. Promoter activity is measured by assaying reporter expression.
25 See, for example, An et al., "Techniques for Isolating and Characterizing Transcription Promoters, Enhancers, and Terminators," in METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY, Glick et al. (eds.), pages 155-166 (CRC Press, 1993).

30 II. Cloning of ALS3 Promoters

A 1063 bp *Xba*I/*Nco*I fragment comprising the promoter region of the ALS3 gene from wild type *Brassica napus* was cloned into pGEM5 (Promega Corporation, Madison, WI) and characterized. The nucleotide sequence of the 1063 bp
35 *Xba*I/*Nco*I fragment was determined by the dideoxy sequencing protocol. Sanger et al., *Proc. Nat'l Acad.*

- 11 -

Sci. USA, 74:5463 (1977). The nucleotide sequence of the 1063 bp *Xba*I/*Nco*I fragment is shown in Figure 1 [SEQ ID NO: 1] and a detailed restriction map of this same fragment is shown in Figure 2.

5 Other ALS3 promoters having substantial sequence similarity with the nucleotide sequence shown in Figure 1 can be cloned by conventional methods. Oligonucleotides of defined sequence are chemically synthesized. Itakura et al., *Annu. Rev. Biochem.* 53:323
10 (1984). Numerous automated and commercially available DNA synthesizers are currently available. The probe can be a single and relatively short oligonucleotide of defined sequence, pools of short oligonucleotides whose sequences are highly degenerate or pools of long
15 oligonucleotides of lesser degeneracy. Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd ed. (Cold Spring Harbor Press, Cold Spring Harbor, N.Y. 1989).

The oligonucleotide hybridization probes based on SEQ ID NO: 1 are labeled, for example radio labeled, by
20 conventional methods and used to detect related nucleotides sequences in *Brassica* genomic libraries by means of DNA hybridization. See, for example, Sambrook *supra*.

A plant genomic DNA library can be prepared by means
25 well-known in the art. See, for example, Slightom et al. "Construction of λ Clone Banks," in *METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY*, Glick et al. (eds.), pages 121-146 (CRC Press, 1993). Genomic DNA can be isolated from *Brassica* tissue, for example, by lysing
30 plant tissue with the detergent Sarkosyl, digesting the lysate with proteinase K, clearing insoluble debris from the lysate by centrifugation, precipitating nucleic acid from the lysate using isopropanol, and purifying resuspended DNA on a cesium chloride density gradient.
35 Ausubel et al. (eds.). *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY*, pages 2.3.1-2.3.3. (1990).

- 12 -

DNA fragments that are suitable for the production of a genomic library can be obtained by the random shearing of genomic DNA or by the partial digestion of genomic DNA with restriction endonucleases. -See, for
5 example, Ausubel et al., *supra*, at pages 5.3.2-5.4.4, and Slightom et al., *supra*.

Genomic DNA fragments can be inserted into a vector, such as a bacteriophage or cosmid vector, in accordance with conventional techniques, such as the use of
10 restriction enzyme digestion to provide appropriate termini, the use of alkaline phosphatase treatment to avoid undesirable joining of DNA molecules, and ligation with appropriate ligases. Techniques for such manipulation are disclosed by Slightom et al., *supra*, and
15 are well-known in the art. Also see Ausubel et al., *supra*, at pages 3.0.5-3.17.5.

A library containing genomic clones is screened with DNA hybridization probes based on the nucleotides sequence of the ALS3 promoter shown in Figure 1 [SEQ ID
20 NO: 1] using standard techniques. See, for example, Ausubel et al., *supra*, at pages 6.0.3-6.6.1; Slightom et al., *supra*.

III. Characterization of ALS3 Promoters

Genomic clones can be analyzed using a variety of
25 techniques such as restriction analysis, Southern analysis, primer extension analysis, and DNA sequence analysis. Primer extension analysis or S1 nuclease protection analysis, for example, can be used to localize the putative start site of transcription of the cloned
30 gene. Ausubel et al., *supra*, at pages 4.8.1-4.8.5; Walmsley et al., "Quantitative and Qualitative Analysis of Exogenous Gene Expression by the S1 Nuclease Protection Assay," in METHODS IN MOLECULAR BIOLOGY, VOL.
7: GENE TRANSFER AND EXPRESSION PROTOCOLS, Murray (ed.),
35 pages 271-281 (Humana Press Inc. 1991). Structural

analysis can be combined with functional analysis for a complete characterization of the promoter region.

5 The general approach of such functional analysis involves subcloning fragments of the genomic clone into an expression vector which contains a reporter gene, introducing the expression vector into various plant tissues, and assaying the tissue to detect the transient expression of the reporter gene. The presence of a constitutive, tissue-general promoter is verified by the
10 observation of reporter gene expression in diverse plant tissues including roots, stems or leaves.

Methods for generating fragments of a genomic clone are well-known. Preferably, enzymatic digestion is used to form nested deletions of genomic DNA fragments. See,
15 for example, Ausubel et al., *supra*, at pages 7.2.1-7.2.20; An et al., *supra*.

Alternatively, DNA that resides "upstream," or 5'-ward, of the transcriptional start site can be tested by subcloning a DNA fragment that contains the upstream
20 region, digesting the DNA fragment in either the 5' to 3' direction or in the 3' to 5' direction to produce nested deletions, and subcloning the small fragments into expression vectors for transient expression.

The selection of an appropriate expression vector
25 will depend upon the method of introducing the expression vector into host cells. Typically, an expression vector contains: (1) prokaryotic DNA elements coding for a bacterial replication origin and an antibiotic resistance marker to provide for the growth and selection of the
30 expression vector in the bacterial host; (2) eukaryotic DNA elements that control initiation of transcription, such as a promoter; (3) DNA elements that control the processing of transcripts, such as a transcription termination/polyadenylation sequence; and (4) a reporter
35 gene that is operably linked to the DNA elements that control transcription initiation. Useful reporter genes

include β -glucuronidase, β -galactosidase, chloramphenicol acetyl transferase, luciferase, and the like. Preferably, the reporter gene is either the β -glucuronidase (GUS) gene or the luciferase gene. See, 5 for example, Jefferson et al., *Plant Molecular Biology Reporter* 5(4):387 (1987). General descriptions of plant expression vectors and reporter genes can be found in Gruber et al., "Vectors for Plant Transformation," in METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY, 10 Glick et al. (eds.), pages 89-119 (CRC Press, 1993). Moreover, GUS expression vectors and GUS gene cassettes are available from Clontech Laboratories, Inc. (Palo Alto, CA), while luciferase expression vectors and luciferase gene cassettes are available from Promega 15 Corporation (Madison, WI).

Expression vectors containing test genomic fragments can be introduced into protoplasts, or into intact tissues or isolated cells. Preferably, expression vectors are introduced into intact tissues. General 20 methods of culturing plant tissues are provided, for example, by Miki et al., "Procedures for Introducing Foreign DNA into Plants," in METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY, Glick et al. (eds.), pages 67-88 (CRC Press, 1993). Methods of introducing expression 25 vectors into plant tissue include the direct infection or co-cultivation of plant tissue with *Agrobacterium tumefaciens*. Horsch et al., *Science* 227:1229 (1985). Descriptions of *Agrobacterium* vector systems and methods for *Agrobacterium*-mediated gene transfer are provided by 30 Gruber et al., *supra*, and Miki et al., *supra*. Methods of introducing expression vectors into plant tissue also include direct gene transfer methods such as microprojectile-mediated delivery, DNA injection, electroporation, and the like. See, for example, Gruber 35 et al., *supra*; Miki et al., *supra*.

- 15 -

The above-described methods have been used to identify and characterize the *Brassica* ALS3 promoter that is constitutively expressed in transformed plants. In particular, the ALS3 constitutive and generally non-tissue-preferred promoter was found to reside within a 1063 bp DNA fragment shown in Figure 1 [SEQ ID NO: 1]. Thus, the present invention encompasses a DNA molecule having a nucleotide sequence of SEQ ID NO: 1 and having the function of a constitutive plant promoter.

Variants of the 1063 bp generally constitutive and non-tissue-preferred promoter can be produced by deleting, adding and/or substituting nucleotides for the nucleotides recited in SEQ ID NO: 1. Such variants can be obtained, for example, by oligonucleotide-directed mutagenesis, linker-scanning mutagenesis, mutagenesis using the polymerase chain reaction, and the like. Ausubel et al., supra, at pages 8.0.3-8.5.9. Also see generally, McPherson (ed.), DIRECTED MUTAGENESIS: A PRACTICAL APPROACH, IRL Press (1991). Thus, the present invention also encompasses DNA molecules comprising nucleotide sequences that have substantial sequence similarity with SEQ ID NO: 1 and function as a tissue-general promoter.

Moreover, additional deletion analyses are performed to further localize the core promoter region within the 1063 bp promoter. Thus, the present invention also encompasses fragments of the DNA molecule having nucleotide sequence of SEQ ID NO: 1, as long as the DNA fragments function as a constitutive promoter.

IV. Vector Construction

The 1063 bp *Xba*I/*Nco*I fragment shown in Figure 1 [SEQ ID NO: 1] was fused to a GUS reporter gene cassette which includes the 3-prime terminator sequence from the potato proteinase inhibitor gene (*PINII*). The ALS3 promoter was cloned as an *Xba*I/*Eco*RI fragment into the corresponding sites of the binary vector pPH11741 to create plasmid

- 16 -

pPHI4960 (Figure 3). Plasmid pPHI1741 differs from plasmid pBI101.1, taught by Jefferson et al., *supra*, by having the CaMV 35S rather than the nopaline synthase 5-prime and 3-prime regulatory sequences driving the NPTII selectable marker gene.

The vector pPHI6333 was constructed by replacing the region between the *PmeI* and *EcoRI* sites of pPHI1741 with a cassette containing the ALS3 promoter driving an NPTII selectable marker gene and the *PIMII* 3' terminator sequence (Figure 4).

V. Agronomic Genes and Selectable Marker Genes for Brassica Transformation

By means of the present invention, agronomic genes and selectable marker genes can be operably linked to the ALS3 promoter and constitutively expressed in transformed plants. More particularly, Brassica can be genetically engineered to express various phenotypes of agronomic interest. The genes implicated in this regard include, but are not limited to, those categorized below.

1. Genes That Confer Resistance To Pests or Disease And That Encode:

- (A) Plant disease resistance genes. Plant defenses are often activated by specific interaction between the product of a disease resistance gene (*R*) in the plant and the product of a corresponding avirulence (*Avr*) gene in the pathogen. A plant variety can be transformed with cloned resistance gene to engineer plants that are resistant to specific pathogen strains. See, for example Jones et al., *Science* 266:789 (1994) (cloning of the tomato *Cf-9* gene for resistance to *Cladosporium fulvum*); Martin et al., *Science* 262:1432 (1993) (tomato *Pto* gene for resistance to *Pseudomonas syringae* pv. tomato encodes a protein kinase); Mindrinos et al., *Cell* 78:1089 (1994)

- 17 -

(*Arabidopsis* RSP2 gene for resistance to *Pseudomonas syringae*).

- 5 (B) A *Bacillus thuringiensis* protein, a derivative thereof or a synthetic polypeptide modeled thereon. See, for example, Geiser et al., *Gene* 48:109 (1986), who disclose the cloning and nucleotide sequence of a Bt δ -endotoxin gene. Moreover, DNA molecules encoding δ -endotoxin genes can be purchased from American Type Culture Collection (Rockville, MD), under ATCC accession Nos. 40098, 67136, 31995 and 31998.
- 10 (C) A lectin. See, for example, the disclosure by Van Damme et al., *Plant Molec. Biol.* 24:825 (1994), who disclose the nucleotide sequences of several *Clivia miniata* mannose-binding lectin genes.
- 15 (D) A vitamin-binding protein such as avidin. See U.S. patent application serial No. 07/911,864, the contents of which are hereby incorporated by reference. The application teaches the use of avidin and avidin homologues as larvicides against insect pests.
- 20 (E) An enzyme inhibitor, for example, a protease inhibitor or an amylase inhibitor. See, for example, Abe et al., *J. Biol. Chem.* 262:16793 (1987) (nucleotide sequence of rice cysteine proteinase inhibitor), Huub et al., *Plant Molec. Biol.* 21:985 (1993) (nucleotide sequence of cDNA encoding tobacco proteinase inhibitor I), and Sumitani et al., *Biosci. Biotech. Biochem.* 57:1243 (1993) (nucleotide sequence of *Streptomyces nitrosporeus* α -amylase inhibitor).
- 25 (F) An insect-specific hormone or pheromone such as an ecdysteroid and juvenile hormone, a variant thereof, a mimetic based thereon, or an antagonist or agonist thereof. See, for
- 30
- 35

- 18 -

example, the disclosure by Hammock et al.,
Nature 344:458 (1990), of baculovirus
expression of cloned juvenile hormone esterase,
an inactivator of juvenile hormone. —

- 5 (G) An insect-specific peptide or neuropeptide
which, upon expression, disrupts the physiology
of the affected pest. For example, see the
disclosures of Regan, J. Biol. Chem. 269:9
10 (1994) (expression cloning yields DNA coding
for insect diuretic hormone receptor), and
Pratt et al., Biochem. Biophys. Res. Comm.
163:1243 (1989) (an allostatin is identified in
Diploptera punctata). See also U.S. patent No.
15 5,266,317 to Tomalski et al., who disclose
genes encoding insect-specific, paralytic
neurotoxins.
- (H) An insect-specific venom produced in nature by
a snake, a wasp, etc. For example, see Pang et
al., Gene 116:165 (1992), for disclosure of
20 heterologous expression in plants of a gene
coding for a scorpion insectotoxic peptide.
- (I) An enzyme responsible for an hyperaccumulation
of a monoterpane, a sesquiterpene, a steroid,
hydroxamic acid, a phenylpropanoid derivative
25 or another non-protein molecule with
insecticidal activity.
- (J) An enzyme involved in the modification,
including the post-translational modification,
of a biologically active molecule; for example,
30 a glycolytic enzyme, a proteolytic enzyme, a
lipolytic enzyme, a nuclease, a cyclase, a
transaminase, an esterase, a hydrolase, a
phosphatase, a kinase, a phosphorylase, a
polymerase, an elastase, a chitinase and a
35 glucanase, whether natural or synthetic. See
PCT application WC 93/02197 in the name of

5 Scott et al., which discloses the nucleotide sequence of a callase gene. DNA molecules which contain chitinase-encoding sequences can be obtained, for example, from the ATCC under
10 accession Nos. 39637 and 67152. See also Kramer et al., *Insect Biochem. Molec. Biol.* 23:691 (1993), who teach the nucleotide sequence of a cDNA encoding tobacco hookworm chitinase, and Kawalleck et al., *Plant Molec. Biol.* 21:673 (1993), who provide the nucleotide sequence of the parsley ubi4-2 polyubiquitin gene.

15 (K) A molecule that stimulates signal transduction. For example, see the disclosure by Botella et al., *Plant Molec. Biol.* 24:757 (1994), of nucleotide sequences for mung bean calmodulin cDNA clones, and Griess et al., *Plant Physiol.* 104:1467 (1994), who provide the nucleotide sequence of a maize calmodulin cDNA clone.

20 (L) A hydrophobic moment peptide. See U.S. patent applications serial No. 08/168,809 (disclosure of peptide derivatives of Tachyplesin which inhibit fungal plant pathogens) and serial No. 08/179,632 (teaches synthetic antimicrobial peptides that confer disease resistance), the
25 respective contents of which are hereby incorporated by reference.

(M) A membrane permease, a channel former or a channel blocker. For example, see the
30 disclosure by Jaynes et al., *Plant Sci.* 89:43 (1993), of heterologous expression of a cecropin- β lytic peptide analog to render transgenic tobacco plants resistant to *Pseudomonas solanacearum*.

35 (N) A viral-invasive protein or a complex toxin derived therefrom. For example, the

- 20 -

- 5 accumulation of viral coat proteins in transformed plant cells imparts resistance to viral infection and/or disease development effected by the virus from which the coat protein gene is derived, as well as by related viruses. See Beachy et al., *Ann. Rev. Phytopathol.* 28:451 (1990). Coat protein-mediated resistance has been conferred upon transformed plants against alfalfa mosaic virus, cucumber mosaic virus, tobacco streak virus, potato virus X, potato virus Y, tobacco etch virus, tobacco rattle virus and tobacco mosaic virus. *Id.*
- 10
- 15 (O) An insect-specific antibody or an immunotoxin derived therefrom. Thus, an antibody targeted to a critical metabolic function in the insect gut would inactivate an affected enzyme, killing the insect. Cf. Taylor et al., Abstract #497, SEVENTH INT'L SYMPOSIUM ON
- 20 MOLECULAR PLANT-MICROBE INTERACTIONS (1994) (enzymatic inactivation in transgenic tobacco via production of single-chain antibody fragments).
- 25 (P) A virus-specific antibody. See, for example, Tavladoraki et al., *Nature* 366:469 (1993), who show that transgenic plants expressing recombinant antibody genes are protected from virus attack.
- 30 (Q) A developmental-arrestive protein produced in nature by a pathogen or a parasite. Thus, fungal endo α -1,4-D-polygalacturonases facilitate fungal colonization and plant nutrient release by solubilizing plant cell wall homo- α -1,4-D-galacturonase. See Lamb et al., *Bio/Technology* 10:1436 (1992). The
- 35 cloning and characterization of a gene which

- 21 -

encodes a bean endopolygalac-turonase-inhibiting protein is described by Toubart et al., *Plant J.* 2:367 (1992).

- (R) A developmental-arrestive protein produced in nature by a plant. For example, Logemann et al., *Bio/Technology* 10:305 (1992), have shown that transgenic plants expressing the barley ribosome-inactivating gene have an increased resistance to fungal disease.

2. Genes That Confer Resistance To A Herbicide, For Example:

- (A) A herbicide that inhibits the growing point or meristem, such as an imidazalinone or a sulfonylurea. Exemplary genes in this category code for mutant ALS and AHAS enzyme as described, for example, by Lee et al., *EMBO J.* 7:1241 (1988), and Miki et al., *Theor. Appl. Genet.* 80:449 (1990), respectively.

- (B) Glyphosate (resistance imparted by mutant EPSP synthase and *aroA* genes, respectively) and other phosphono compounds such as glufosinate (PAT and *bar* genes), and pyridinoxy or phenoxy proprionic acids and cyclohexones (ACCase inhibitor-encoding genes). See, for example, U.S. patent No. 4,940,835 to Shah et al., which discloses the nucleotide sequence of a form of EPSP which can confer glyphosate resistance. A DNA molecule encoding a mutant *aroA* gene can be obtained under ATCC accession No. 39256, and the nucleotide sequence of the mutant gene is disclosed in U.S. patent No. 4,769,061 to Comai. European patent application No. 0 333 033 to Kumada et al. and U.S. patent No. 4,975,374 to Goodman et al. disclose nucleotide sequences of glutamine synthetase genes which

- 22 -

- confer resistance to herbicides such as L-phosphinothricin. The nucleotide sequence of a phosphinothricin-acetyl-transferase gene is provided in European application No. 0 242 246 to Leemans et al. De Greef et al., *Bio/Technology* 7:61 (1989), describe the production of transgenic plants that express chimeric bar genes coding for phosphinothricin acetyl transferase activity. Exemplary of genes conferring resistance to phenoxy proprionic acids and cyclohexones, such as sethoxydim and haloxyfop, are the *Acc1-S1*, *Acc1-S2* and *Acc1-S3* genes described by Marshall et al., *Theor. Appl. Genet.* 83:435 (1992).
- 15 (C) A herbicide that inhibits photosynthesis, such as a triazine (*psbA* and *gs+* genes) and a benzonitrile (nitrilase gene). Przibilla et al., *Plant Cell* 3:169 (1991), describe the transformation of *Chlamydomonas* with plasmids encoding mutant *psbA* genes. Nucleotide sequences for nitrilase genes are disclosed in U.S. patent No. 4,810,648 to Stalker, and DNA molecules containing these genes are available under ATCC accession Nos. 53435, 67441 and 67442. Cloning and expression of DNA coding for a glutathione S-transferase is described by Hayes et al., *Biochem. J.* 285:173 (1992).
- 25
3. Genes That Confer Or Contribute To A Value-Added Trait, Such As:
- 30 (A) Modified fatty acid metabolism, for example, by transforming maize *Brassica* with an antisense gene of stearyl-ACP desaturase to increase stearic acid content of the plant. See Knultzon et al., *Proc. Nat'l Acad Sci. USA* 89:2624 (1992).
- 35

(B) Decreased phytate content

- 5 (1) Introduction of a phytase-encoding gene would enhance breakdown of phytate, adding more free phosphate to the transformed plant. For example, see Van Hartingsveldt et al., *Gene* 127:87 (1993), for a disclosure of the nucleotide sequence of an *Aspergillus niger* phytase gene.
- 10 (2) A gene could be introduced that reduces phytate content. In maize, this, for example, could be accomplished, by cloning and then re-introducing DNA associated with the single allele which is responsible for maize mutants
- 15 characterized by low levels of phytic acid. See Raboy et al., *Maydica* 35:383 (1990).

- 20 (C) Modified carbohydrate composition effected, for example, by transforming plants with a gene coding for an enzyme that alters the branching pattern of starch. See Shiroza et al., *J. Bacteriol.* 170:810 (1988) (nucleotide sequence of *Streptococcus mutans* fructosyltransferase gene), Steinmetz et al., *Mol. Gen. Genet.* 25 200:220 (1985) (nucleotide sequence of *Bacillus subtilis* levansucrase gene), Pen et al., *Bio/Technology* 10:292 (1992) (production of transgenic plants that express *Bacillus licheniformis* α -amylase), Elliot et al., *Plant Molec. Biol.* 21:515 (1993) (nucleotide sequences of tomato invertase genes), Søgaard et al., *J. Biol. Chem.* 268:22480 (1993) (site-directed mutagenesis of barley amylase gene), and Fisher et al., *Plant Physiol.* 102:1045 35 (1993) (maize endosperm starch branching enzyme II).

4. Selectable Marker Genes:

(A) Numerous selectable marker genes are available for use in plant transformation including, but not limited to, neomycin phosphotransferase I, hygromycin phosphotransferase, EPSP synthase and dihydropteroate. See Miki et al., "Procedures for Introducing Foreign DNA into Plants," in METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY, Glick et al. (eds.), pages 67-88 (CRC Press, 1993).

Synthesis of genes suitably employed in the present invention can be effected by means of mutually priming, long oligonucleotides. See, for example, Ausubel et al. (eds.), CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, pages 8.2.8 to 8.2.13 (Wiley Interscience 1990), and Wosnick et al., Gene 60:115 (1987). Moreover, current techniques which employ the polymerase chain reaction permit the synthesis of genes as large as 1.8 kilobases in length. See Adang et al., Plant Molec. Biol. 21:1131 (1993), and Bambot et al., PCR Methods and Applications 2:266 (1993).

The present invention, thus generally described, will be understood more readily by reference to the following examples, which are provided by way of illustration and are not intended to be limiting of the present invention.

Example 1

Plant Transformation and Expression Analysis

The vector pPHI4960, shown in Figure 3, was introduced into the *Agrobacterium* strain GV3101 by transformation. *B. napus* cultivar Westar was used throughout this experiment. Transgenic *B. napus* plants were generated by *Agrobacterium* co-cultivation of cotyledonary petioles and microspore derived embryos. See Arnoldo et al., Genome 35:58 (1992). Parallel transformations were carried out using pPHI3042 in order to compare the strength of the ALS3 promoter with that of

- 25 -

the CaMV 35S promoter. Plasmid pPHI3042 is shown in Figure 5 and contains the CaMV 35S promoter operably linked to the GUS gene. Tissues from plants containing independent transformation events were quantitatively
5 analyzed for GUS expression using a histochemical assay method based on that of Jefferson R.A., *Plant Mol. Biol. Rep.* 5:387 (1987).

Table 1 summarizes quantitative GUS expression data from *B. napus* plants independently transformed with
10 plasmid pPHI4960 (HWSC63E, HWSC63F, HWSC63G, HWSC63H, PWSC174C, PWSC174D, PWSC174G, PWSC174J, PWSC174M, PWSC174N, and PWSC176A). Tissue from the primary transgenic plant (T0) was analyzed. Plasmid pPHI4960 is
15 shown in Figure 3 and contains the *B. napus* ALS3 promoter operably linked to the GUS gene. The controls consisted of the same nontransformed cultivar 'Westar' and 4 independent *B. napus* plants transformed with the plasmid pPHI3042 (P105B, P105E, P126B and P145).

Table 1
SUMMARY OF BRASSICA NAPUS TRANSFORMED
WITH pPHI4960 AND pPHI3042

Transgenic	GUS Expression (T0) Fluorogenic (pmoles/MU/h/ μ g protein)		Southern PINII Probe
	leaf	root	
HWSC63E ¹	0.2	6.2	5
HWSC63F ¹	131	167	1
HWSC63G ¹	416	246	1
HWSC63H ¹	21	29	1
PWSC174C ¹	7.5	63.6	1
PWSC174D ¹	11	17	1
PWSC174G ¹	309	no root	0/1
PWSC174J ¹			1
PWSC174M ¹			0/1
PWSC174N ¹			1
PWSC176A ¹	0.2	37	4
P105B ²	316	143	
P105E ²	241	279	
P126B ²	0.4	1.2	
P145 ²	0.2	0.6	
WESTAR ³	0.1	11.2	

¹ *B. napus* transformed with pPHI4960

² *B. napus* transformed with pPHI3042

³ Non-transgenic control

The data reveal that there is variability in the level of expression in the transformants. Variability in the level of expression of transformed genes has been noted in numerous transformation experiments described in the literature and has been attributed to such factors as differences in copy number, position effects and co-suppression. However, plants expressing high levels, or a desired level, of the foreign protein can be identified and selected using the routine screening methods of the present invention.

The data also reveal that there are essentially two classes of expression levels in plants transformed with plasmid pPHI4960. The first class consists of transformed plants expressing high levels of assayable product (131-416 pmol of methylumbelliferone (MU)/h/ μ g protein). The second class consists of transformed plants expressing low levels of assayable product (0.2-37 pmol MU/h/ μ g protein). The low expression level is generally comparable to the non-transformed control plants examined. These values do not imply absolute expected ranges but represent the maximum and minimum values for each class in the present study.

Southern blot analysis of total genomic DNA isolated from transformed plants was undertaken to determine the copy number of the GUS gene cassette in each transformant. The *PINII* terminator was used as the radiolabeled hybridization probe. Total genomic DNA was isolated from each plant transformant using a variation of a CTAB protocol. See, for example, Dellaporta et al., 1983, *Plant Mol. Biol. Rep.* 1(4):19-21, or Saghai-Maroot et al., 1984, *PNAS* 81:8014-8018. The total genomic DNA was digested with a restriction enzyme(s), separated on agarose gels by means of electrophoresis and hybridized with the radiolabeled *PINII* terminator sequence as a probe. Plasmid pPHI4960 was digested with either *HindIII* alone to assess number of integrations, or double digested with *HindIII* plus *EcoRI* to determine copy number. Similarly, pPHI3042 was digested with *HindIII*

alone to assess number of integrations, or with EcoRI alone to evaluate copy number (see enzyme cut sites on maps in Figures 3 and 5, respectively). Although an exact copy number for each integration event was not
5 determined, the relative intensities of hybridizing bands were used to compare Southern data between transgenics with different vectors.

Plants transformed with multiple copies of the plasmid pPHI4960 (HWSC63E and PWSC176A) produced low
10 levels of MU. On the other hand, 50% of the single integration events produced high levels of MU. Plants with single transformation events produced high levels of MU irrespective of whether the GUS gene was expressed from the CaMV 35S or the ALS3 promoter.

15 Among the transformed plants producing high levels of MU, plants transformed with the ALS3 GUS cassette produced just as much MU as plants transformed with the CaMV 35S GUS cassette, irrespective of whether enzyme activity was measured in the roots or shoots. Since the
20 CaMV 35S promoter is regarded as a strong constitutive, generally non-tissue-preferred promoter, with wide application in expressing genes in transgenic plants, the Brassica ALS3 promoter offers a suitable alternative to the CaMV 35S promoter.

25 The level of GUS expression measured in leaf tissue of HWSC63G was higher than the GUS level produced in leaf or root tissue of plants transformed with the CaMV 35S GUS cassette. Accordingly, the ALS3 promoter may be more effective than the CaMV 35S promoter for expression of
30 certain genes at high levels. For example, the ALS3 promoter may be preferable to the CaMV 35S promoter in those instances where high level expression of the transformed gene is required. It is understood by those skilled in the art that the range of expression in
35 transgenic plants can vary among transformation events. The present data indicates that it is possible to recover at least some events with activity comparable to that

obtained with the 35S promoter by means of routine screening using the methods disclosed herein.

To test the usefulness of the ALS3 promoter for expression of operably linked genes in other species, transformations were also conducted in sunflower. Sunflower leaf discs were co-cultivated with *Agrobacterium* carrying either pPHI4960 or pPHI6333. See Malone-Schoneberg, et al., 1994, *Plant Science* 103: 199-207. Stably transformed calli were selected using 100 mg/L of kanamycin. GUS or NPTII expression levels were determined for a number of independent transformation events, using histochemical staining for GUS expression analysis, and semi-quantitative ELISAs for NPTII expression analysis. While GUS expression varied among individual plants, the ranges of expression from the two promoters in leaf tissues overlapped substantially (73-3119 fluorescence units for the ALS3 promoter and 40-2788 fluorescence units for the 35S promoter). Limited ELISAs also yielded comparable results for NPTII expression from the two promoters (data not shown), although extensive whole plant analysis was not carried out.

Example 2

Activity of the ALS3 Promoter in Transgenic B. napus in Selected Tissues from Seedling, Vegetative, Flowering and Mature Stage Plants

B. napus cv. Westar was transformed with pPHI4960 as described in Example 1. Four independent transformants were analyzed (HWSC63F, HWSC63G, HWSC63E and PWSC176A). The primary transformants are designated T0. The seed produced by T0 plants are T1 seeds and the plants produced by T1 seeds are referred to as T1 plants.

GUS expression was quantitatively determined, by the method described in Example 1, in T1 plants at the seedling stage (less than 2 true leaves), the vegetative stage (4-5 true leaves), the early flowering stage (about 20-80% full bloom), and in mature plants. In addition, GUS expression was quantitatively determined in specific plant tissues, including meristem, cotyledon, hypocotyl,

leaf, petiole, stem, root, pod, seed and flowers. "Meristem" refers to the meristematic apex and includes much of the tissue surrounding the true meristem. "Flowers" refers to all organs associated with the flowers, including ovaries, anthers and petals. "Pods" refers to the developing green pods containing green and brown seed. "Seed" refers to the seed still on the plant, hence the seeds were not desiccated. The results of quantitative enzyme assays were confirmed by means of histochemical assays as in example 1.

GUS expression in *B. napus* cv. Westar transformed with pHI4960 was compared to plants of the same cultivar transformed with pPHI3042. A non-transgenic Westar line was used as a control. Although T1 stage plants are segregating with respect to the transformed genes, only T1 plants that had a transgene, as evidenced by expression of neomycin phosphotransferase or survival on kanamycin, were analyzed. Seeds were sown on an agar plate containing 100 µg/ml of kanamycin. Those seeds which germinated and resulted in green seedlings were selected as transformed segregants. Those which bleached white were non-transformed segregants and were discarded or used as negative controls.

Table 2 summarizes Southern, expression and segregation data for several transgenics. GUS expression was measured in leaf tissue from plants at the 3-7 leaf stage. In those primary transgenics with a simple integration pattern, which either showed no segregation (and were therefore fixed) or whose subsequent progeny segregated in approximately a 3:1 ratio with respect to the transformed genes, the levels of GUS expression from both the 35S and ALS3 promoters was relatively high. In contrast, those with complex integration patterns and/or with complex segregation had low GUS expression.

Table 2
ALS Promoter Study

	Southern (T0)	T0 GUS assay results	T1 GUS assay results	Segregation
35S Transgenics				
NWSC49J	8 INSERTS	LOW	LOW	23:1
NWSC49B	1 INSERT	HIGH	HIGH	3:1 (18:6)
NWSC49F	1 INSERT	MEDIUM	LOW	3:4
dhNWSC49G	2 INSERTS	MEDIUM	HIGH/MED	14:0 (FIXED)
ALS Transgenics				
PWSC176A	4 INSERTS	MEDIUM	HIGH	4:1 (16:4)
HWSC63E	5 INSERTS	LOW	LOW	2:1 (9:5)
HWSC63F	1 INSERT	HIGH	MED/HIGH	7:1 (14:2)
HWSC63G	1 INSERT	HIGH	HIGH	7:0

Figures 7, 8, 9 and 10 present GUS expression data in specific tissues taken from plants at the seedling, vegetative, flowering and mature stage, respectively. As evidenced by the results shown in Figure 7, the ALS3 promoter functions in seedlings, specifically in cotyledons, hypocotyls and "meristematic apex." The ALS3 promoter drives low level expression of GUS in the roots at the seedling stage. At the 4-5 true leaf stage, the ALS promoter functions in leaves, petioles, stem, "meristematic apex" and roots, as shown in Figure 8. At the flowering stage, the ALS3 promoter functions in flowers, leaves, stem and roots (Figure 9). Histochemical analysis of flowers revealed expression of GUS in petals, stigma, anthers and pollen. GUS activity was detected in all tissues (leaf, stem pod and seed) taken from mature *B. napus* plants transformed with pPHI4960. Accordingly, the ALS3 promoter is constitutively expressed in diverse plant tissues throughout the development of the plant.

20

*Example 3**Selectable Marker Genes Expressed
from the ALS3 Promoter*

The ALS3 promoter can be used to drive expression of selectable marker genes for plant transformation. *B. napus* was transformed with plasmids pPHI1741, pPHI3042, pPHI5765 and pPHI6333 (Figure 4) using the methods described in Example 1. Transformants were selected on medium containing 100 µg/ml kanamycin. Plasmid pPHI1741 contains the CaMV 35S promoter operably linked to the NPTII structural gene and the terminator sequence of the 195 gene. Plasmid pPHI3042 contains two expression cassettes with the CaMV 35S promoter operably linked to the NPTII structural gene and the CaMV 35S promoter operably linked to the GUS gene. Plasmid pPHI5765 contains the CaMV 35S promoter operably linked to the NPTII structural gene. Finally, plasmid pPHI6333, shown in Figure 4, contains the ALS3 promoter operably linked

to the *NPTII* structural gene and the terminator sequence of the potato proteinase inhibitor gene (*PINII*).

Table 3
Selectable Marker Genes Expressed
From the ALS3 Promoter

VECTOR	TOTAL NUMBER OF EXPLANTS COCULTIVATED	TOTAL NUMBER OF POSITIVES	PERCENT TRANS- FORMED
pPHI1741	1350	64	4.7%
pPHI3042	350	2	0.9%
pPHI5765	1935	60	3.1%
pPHI6333	1935	12	0.6%

As shown in Table 3, the ALS3 promoter is effective in driving expression of the *NPTII* selective marker gene in canola. Transformation of plasmids pPHI1741, pPH3042 and pPHI5765 resulted in the recovery of 4.7%, 0.9% and 3.1% transformants among the total *B. napus* explants cocultivated with these plasmids, respectively. A total of 12 transformants were obtained from experiments in which pPHI6333 was cocultivated with 1,935 *B. napus* explants, or 0.6% of the explants were transformed. Ten of the 12 transformants expressed low levels of *NPTII*. In sunflower, pPHI6333 was as effective as pPHI4960 for selection of transformed plants; that is, both the 35S and the ALS3 promoter allowed selection of transformed tissues at 100 mg/L of kanamycin. Accordingly, the ALS3 promoter is effective in driving expression of a selective marker gene for plant transformation.

Although the foregoing refers to particular preferred embodiments, it will be understood that the

present invention is not so limited. It will occur to those of ordinary skill in the art that various modifications may be made to the disclosed embodiments and that such modifications are intended to be within the scope of the present invention, which is defined by the following claims.

All publications and patent applications mentioned in this specification are indicative of the level of skill of those in the art to which the invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference in its entirety.

What Is Claimed Is:

1. An isolated DNA molecule comprising a promoter operably linked to a foreign structural gene, wherein said promoter comprises a nucleotide sequence
5 corresponding to the sequence of a polynucleotide from the group consisting of (a) a *XbaI/NcoI* fragment 5-prime to the *Brassica napus* ALS3 structural gene or (b) a nucleotide sequence that has substantial sequence similarity with said *XbaI/NcoI* fragment.
- 10 2. An isolated DNA molecule comprising a promoter operably linked to a foreign structural gene, wherein said promoter comprises a nucleotide sequence corresponding to the sequence of a polynucleotide from the group consisting of (a) SEQ ID NO: 1 or (b) a
15 nucleotide sequence that has substantial sequence similarity with SEQ ID NO: 1.
3. The DNA molecule of claims 1 or 2, wherein said foreign structural gene is an agronomically important gene or a selectable marker gene.
- 20 4. An expression vector comprising the isolated DNA molecule of claim 1.
5. An expression vector comprising the isolated DNA molecule of claim 2.
- 25 6. A transformed host comprising the expression vector of claim 4.
7. The transformed host comprising an expression vector of claim 5.
8. A method of using an ALS3 promoter to produce a foreign protein in a transformed host plant, comprising
30 the steps of:
 - (a) constructing an expression vector comprising a promoter operably linked to a foreign structural gene, wherein said promoter comprises a nucleotide sequence corresponding to the sequence of a polynucleotide from
35 the group consisting of (i) a *XbaI/NcoI* fragment 5-prime to the *Brassica napus* ALS3 structural gene or (ii) a

nucleotide sequence that has substantial sequence similarity with said *XbaI/NcoI* fragment; and

(b) transforming a host.

5 9. A method of using an ALS3 promoter to produce a foreign protein in a transformed host plant, comprising the steps of:

10 (a) constructing an expression vector comprising a promoter operably linked to a foreign structural gene, wherein said promoter comprises a nucleotide sequence corresponding to the sequence of a polynucleotide from the group consisting of (i) SEQ ID NO: 1 or (ii) a nucleotide sequence that has substantial sequence similarity with SEQ ID NO: 1; and

(b) transforming a host.

FIG. 1

```

-1059  ICTAGATTAA CATATGCGAC CACTCAGGTT AATAATTCTC CCAAATAACA
          XbaI
-1009  TACATTGGTA TGAAGATTAT GTCTAAATCA AACATAAAAT CGACTCCAAA
-959  TACGAACCCA CAAACTTACA AGTAAATCAA ACATAAAATT TATCTGCATC
-909  CGACTATGTC TACGCCATTG TTTCACCACT CACCAATTAT AGAGGGACAA
-859  ACCTACAGAT CATACTATAT CTAACAGATT TCAACCCCTA AGCAATTCAC
-809  AGAATTGACA TGGACATAGC TAGAGCAAAC CTGCATGATG AAAGAAGGGC
-759  ACCCACAGTG GCGGAAAAG ATGAACTTT GACAATACAC ATTGACAAGA
-709  TTGTATGAAG TGGGGTTATA AATTCA TTGC AATGCAATCA TATCTATCAT
-659  CACCAGTTCA TAA TTGACTG GATCCGAATG AACATAAGAT GCTCTACATG
-609  TCGAGACACA ATACTTTAGT CATAATGTTT CTCATTATCA AGTACAGGTA
-559  TTTCTACTTT AAAC TTATAT TATAAGCAAG TATTACAAAT AATGAAATAA
-509  AGACTATGTT TTCTGTTTT AGATGTTTTG ATCTTCATAT TTAAACCAAT
-459  CGTCTTCATT ATTGCAAGAA ACACAATTTT CTAACCTCTG TAACACTCCT
-409  TATAAAATA TATGACATAT GAAACGGAGC GGAGGCTTGC AGTAGACGTA
-359  ACAACATTTT AAAATTGAAC AAAGTATTGC TTAAAAAATA GAATTTGAAG
-309  GCCTTCTAAG CTGAACATAA AATAAAATTA ACACAAAATT ATATCTTTCA
-259  TCACAACCAA CTGAACATAA AATAAAATTA ACACAAAATT ATATCTTTCA
-209  TAAAAACCGA TACATCAAAT TCCGCGCGTA GCGGGACCC TCCCTAGTAA
-159  TTAATACAGT AAAGAAAAGA CCAACACAAAC AAAAATCATA TTCCAAGGGT
-109  ATTTTCGTAA ACAAAACAAA CCCTCACAAAG CCTCGTTTTA TAAAAACGAT
-59  TCACGTTTCA AACTCATTC ATCATCTCTC TCTCATTTCT CTCTCTCTCT
          +1
-9  CATCTAACCAITGG NcoI

```


2/10

XbaI NdeI DdeI MslI
 TCTAGATTAAACATATGCGACCACTCAGGTTAATAATTCTCCCAAATAACATACATTGGTATGAAGATTA
 TGTCTAAATCAAACATAAAATCGACTCCAAATACGAACCCACAAACTTACAAGTAAATCAAACATAAAA
 FokI AspI AccI
 TTTATCTGCATCCGACTATGTCTACGCCATTGTTTCACCACTCACCAATTATAGAGGGACAAACCTACA
 DpnI DdeI AluI
 GATCATACTATATCTAACAGATTTCAAACCCCTAAGCAATTACAGAAATTGACATGGACATAGCTAGAGC
 BspMI BanI
 AAACCTGCATGATGAAAGAGGGCACCCACAGTGGCGGAAAAAGATGAACTTTTGACAAATACACATTGA
 MslI
 CAAGATTGTATGAAGTGGGTTATAAATTTCATTGCAATGCAATCATATCTATCATCACCAGTTCATAAT
 DpnI
 BamHI AflIII
 BstI NspI
 TGACTGGATCCGAATGAACATAAGATGCTCTACATGTCGAGACACAATACTTTAGTCATAATGTTTCTC
 RsaI DraI
 ATTATCAAGTACAGGTATTTCTACTTTTAAACTTATATTATAAGCAAGTATTACAAAATAATGAAATAAAG
 DpnI BbsI
 ACTAATGTTTCCTGTTTAGATGTTTCGATCTTCATATTTTAAACCAATCGTCTTCATTTATGCAAGAAA
 NdeI
 CACAATTTCCCTAACTCCTGTAACACTCCTTATAAAAATATATGACATATGAAACGGAGCGGAGCTTGC
 HaeIII
 StuI
 AccI DraI XmnI DdeI
 AGTAGACGTAAACAACATTTTAAAAATTGAACAAAGTATTGCTTAAAAATAGAAATTTGAAGGCCTTCTAA
 GTTTTAATAAAATTTAACACTAATCACTTAAATATATCACCTCACAACTGAACATATAAAATAAAAT
 AsuI
 TAAACAAAAATTATATCTTTTCATAAAAAACCGATACATCAAAATTCGCGGTAGCGGACCCCTCCCTAG
 HhaI HhaI AvalI
 AsnI Styl
 TAAATTAATACAGTAAAGAAAAGACCACCAACAAACAAATAATCATATTTCCCAAGGGTATTTTCGTAAACAAAC
 Tfil
 AAAACCCCTCACAAGCCTCGTTTTTATAAAAAACGATTTCAGTTTCACAAACTCATTCATCTCTCTCTCA
 NcoI
 Styl
 TTTCTCTCTCTCTCTCTCATCTAACCATGG

FIG. 2


3/10

FIG. 3




4/10

FIG. 4




5/10

FIG. 5




6/10


FIG. 6



7/10




8/10


FIG. 8 VEGETATIVE STAGE GUS ACTIVITY WITH 35S AND ALS PROMOTERS

9/10

FIG. 9 FLOWERING STAGE GUS ACTIVITY WITH 35S AND ALS PROMOTERS



10/10



INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/US 96/03635

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/82 C12N15/60 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MOL GEN GENET 229 (1). 1991. 31-40., XP002008880 RUTLEDGE R G ET AL: "MOLECULAR CHARACTERIZATION AND GENETIC ORIGIN OF THE BRASSICA-NAPUS ACETOHYDROXY ACID SYNTHASE MULTIGENE FAMILY."	1
Y	see the whole document ---	2-9
X	PLANT J. (1992), 2(3), 321-30 , XP002008881 OUELLET, THERESE ET AL: "Members of the acetoxyacid synthase multigene family of Brassica napus have divergent patterns of expression" see the whole document ---	1
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *A* document member of the same patent family

Date of the actual completion of the international search

19 July 1996

Date of mailing of the international search report

29.07.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Maddox, A

INTERNATIONAL SEARCH REPORT

Int. onal Application No
PCT/US 96/03635

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	PLANT PHYSIOLOGY, vol. 91, 1989, pages 574-580, XP002008882 GABARD, J.M., ET AL.: "Cross-resistance to short residual sulfonylurea herbicides in transgenic tobacco plants" see the whole document ---	2-9
P,Y	WO,A,96 07746 (AMERICAN CYANAMID CO ;DIETRICH GABRIELE (NL); SMITH JANE (US); PEN) 14 March 1996 see the whole document ---	2-9
A	MOLECULAR AND GENERAL GENETICS, vol. 219, no. 3, 1 November 1989, pages 413-420, XP000095145 WIERSMA P A ET AL: "ISOLATION, EXPRESSION AND PHYLOGENETIC INHERITANCE OF AN ACETOLACTATE SYNTHASE GENE FROM BRASSICA NAPUS" see the whole document ---	1-9
A	THEOR APPL GENET 80 (4). 1990. 449-458., XP000575963 MIKI B L ET AL: "TRANSFORMATION OF BRASSICA-NAPUS CANOLA CULTIVARS WITH ARABIDOPSIS-THALIANA ACETOHYDROXY ACID SYNTHASE GENES AND ANALYSIS OF HERBICIDE RESISTANCE." see the whole document ---	1-9
A	KEYSTONE SYMPOSIUM ON CROP IMPROVEMENT VIA BIOTECHNOLOGY: AN INTERNATIONAL PERSPECTIVE, KEYSTONE, COLORADO, USA, APRIL 10-16, 1992. J CELL BIOCHEM SUPPL 0 (16 PART F). 1992. 204., XP002008884 BARBOUR E ET AL: "ISOLATION AND CHARACTERIZATION OF A BRASSICA-NAPUS MUTANT ALS GENE AND EXPRESSION IN TRANSGENIC PLANTS." see abstract Y103 ---	1-9
A	MOLECULAR AND GENERAL GENETICS, vol. 246, 20 February 1995, pages 419-425, XP002008914 HATTORI, J., ET AL.: "An acetohydroxy acid synthase mutant reveals a single site involved in multiple herbicide resistance" see page 420, right-hand column ---	1-9

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/03635

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>CROP SCIENCE, vol. 34, 1994, pages 226-229, XP000576642 BRANDLE, J.E., ET AL.: "A comparison of two genes for sulfonylurea herbicide resistance in transgenic tobacco seedlings" see the whole document ----</p>	1-9
A	<p>EP,A,0 559 603 (SANDOZ LTD ;SANDOZ AG (DE); SANDOZ AG (AT)) 8 September 1993 see the whole document -----</p>	1-9

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/03635

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9607746	14-03-96	AU-B- 3544895	27-03-96
EP-A-0559603	08-09-93	AU-B- 661359	20-07-95
		AU-B- 3110493	15-07-93
		CA-A- 2086863	10-07-93
		CN-A- 1079779	22-12-93
		CZ-A- 9203923	16-03-94
		HU-A- 69949	28-09-95
		JP-A- 5276951	26-10-93
		SK-A- 392392	09-08-95
		ZA-A- 9300145	08-07-94